



Published in final edited form as:

*Health Educ Behav.* 2014 October ; 41(10): 98S–107S. doi:10.1177/1090198114540462.

## Health-Related Quality of Life and Health Behaviors in a Population-Based Sample of Older, Foreign-Born, Chinese American Adults Living in New York City

Laura C. Wyatt, MPH<sup>1</sup>, Dr Chau Trinh-Shevrin, PH<sup>1</sup>, Nadia S. Islam, PhD<sup>1</sup>, and Dr Simona C. Kwon, PH<sup>1</sup>

<sup>1</sup>New York University School of Medicine, New York, NY, USA

### Abstract

Although the New York City (NYC) Chinese population aged ≥ 65 increased by 50% between 2000 and 2010, the health needs of this population are poorly understood. Approximately 3,001 Chinese individuals from high density Asian American NYC areas were included in the REACH U.S. Risk Factor Survey; 805 (26.8%) were aged ≥ 65 and foreign-born. Four health-related quality of life (HRQOL) and three behavioral risk factor outcome variables were examined. Descriptive statistics were conducted by gender, and logistic regression models assessed socio-demographic and health factors associated with each outcome. Few women were current smokers (1.3% vs. 14.8% of men), 19% of respondents ate fruits and vegetables ≥ 5 times daily, and one third of individuals received sufficient weekly physical activity. Days of poor health were similar to the national population aged ≥ 65, while self-reported fair or poor health was much greater among our Chinese sample; over 60% of respondents rated their health as fair or poor. Lower education and lower obesity were significantly associated with cigarette smoking among men, and older age was significantly associated with insufficient physical activity overall. Female gender was significantly associated with all poor health days; older age was significantly associated with poor days of physical health, and lower income was significantly associated with poor days of physical health and fair or poor self-reported health. This study provides important health-related information on a rapidly growing older population and highlights future research areas to inform culturally-appropriate health promotion and disease prevention strategies and policies within community-based settings.

### Keywords

community health; health behavior; immigration health; mental health; physical activity; Asian/Pacific Islander; smoking and tobacco use; nutrition

**Corresponding Author:** Laura C. Wyatt, New York University School of Medicine, 550 First Ave, VZN 838F, New York, NY 10016, USA..

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Introduction

Chinese Americans are the largest Asian American subgroup, representing nearly one-quarter (22.8%) of the Asian population in the United States (U.S.) and almost half (47.0%) of the Asian population in New York City (NYC) (U.S. Census Bureau, 2010). Chinese Americans aged 65 make up approximately 11.8% of the NYC Chinese population and experienced a 49.8% growth between 2000 and 2010; conversely, the overall population aged 65 in NYC increased 5.9% during this same period (U.S. Census Bureau, 2000, 2010). European-born older populations are more likely to have entered the U.S. prior to 1970, whereas Asian-born older populations began entering following the Immigration Act of 1965, with an influx in the 1970s (He & U.S. Census Bureau, 2002; Trinh-Shevrin, Islam, & Rey, 2009).

Older Chinese Americans in NYC have been overlooked by health providers and policy makers, as data on this rapidly increasing subgroup are often aggregated with other Asian groups, masking possible differences in health. Research suggests that older Asian Americans are a heterogeneous group exhibiting different characteristics from the general aging population (Trinh-Shevrin et al., 2009). For example, older NYC Chinese individuals live in larger households, have lower incomes, and experience more English language limitations than other groups (Asian American Federation of New York, 2003). Local and national surveys of Asian Americans have included few Asian languages, provide limited geographic coverage, and are based on small sample sizes (Islam et al., 2010). Limited research exists on the health of older Chinese Americans, especially in NYC (G. Kim, Chiriboga, et al., 2010; Parikh, Fahs, Shelley, & Yerneni, 2009; Wong, Dixon, Gilbride, Kwan, & Stein, 2013). Although data from the NYC Community Health Survey exists, information is generally aggregated across Asian Americans, Native Hawaiians and Pacific Islanders and results limited by small sample size (New York City Department of Health and Mental Hygiene, 2012).

Studies focusing on older Chinese Americans reveal this group is at risk for various health conditions, including chronic diseases and poor mental health, which increase with age (Hwang, Chun, Takeuchi, Myers, & Siddarth, 2005; Lau et al., 2005; Rajpathak & Wylie-Rosett, 2011; Shah et al., 2010). Modifiable behavioral risk factors, including inadequate levels of physical activity (PA), smoking, and poor nutrition have an adverse effect on health (Danaei et al., 2009; Mozaffarian et al., 2012), as well as on chronic diseases associated with the leading causes of morbidity and mortality (Buttar, Li, & Ravi, 2005; Lloyd-Jones et al., 2006; Paganini-Hill, 2011; Physical Activity Guidelines Advisory Committee, 2008). A healthy lifestyle contributes to healthy aging (Haveman-Nies, de Groot, & van Staveren, 2003).

Good physical and mental health are important for older adults (Drewnowski & Evans, 2001; McGuire, Strine, Okoro, Ahluwalia, & Ford, 2007). PA helps reduce risk of falls and fractures, as well as the risk of depression (Gregg, Pereira, & Caspersen, 2000; Strawbridge, Deleger, Roberts, & Kaplan, 2002). According to the 1998-2008 National Health Interview Survey, one-third of people aged 65 reported sufficient weekly PA (Carlson, Fulton, Schoenborn, & Loustalot, 2010). Risk of poor mental health varies among different

subgroups of Asian Americans. Filipino American women are shown to have better self-reported health than Chinese and Vietnamese American women (Appel, Huang, Ai, & Lin, 2011); while rates of depression have been found to be high among Chinese Americans, and lower among Filipinos and Vietnamese (Takeuchi et al., 2007; Yeung et al., 2004). Mental health issues, however, are often under-recognized and under-treated among these Asian subgroups (Alegria et al., 2008; G. Kim, Jang, Chiriboga, Ma, & Schonfeld, 2010; Yeung, Yu, Fung, Vorono, & Fava, 2006).

Data for this study were obtained from the Racial and Ethnic Approaches to Community Health across the U.S. (REACH U.S.) Risk Factor Survey. The objectives of the current study were to: (a) describe the demographic and health characteristics of foreign-born, Chinese Americans aged 65 living in NYC; (b) examine health related quality of life (HRQOL) and health behaviors (i.e., PA, fruit and vegetable intake [F&V intake], and smoking); and (c) examine the factors associated with HRQOL and health behaviors. This study fills an important gap in the knowledgebase about the health of this growing population of older Chinese individuals in the U.S., and specifically in NYC.

## Methods

### Study Setting

REACH U.S., supported by the Centers for Disease Control and Prevention (CDC), is designed to mobilize local communities to implement and evaluate community-based strategies to eliminate health disparities within racial and ethnic populations. Surveys were conducted annually in 28 grantee communities as part of evaluation activities. This study used 4 years of aggregated data (2009-2012) from the B Free CEED: National Center of Excellence in the Elimination of Hepatitis B Disparities program site (Pollack et al., 2011; Trinh-Shevrin et al., 2011) that oversampled Asian Americans in NYC.

### Study Design and Sample

The survey used an address-based sampling design with geographical information systems technology to target Census tracts with a large density of Asian Americans, and was conducted using self-administered questionnaires (60%), telephone interviews (39%), and in-person interviews (1%, 2009). Chinese respondents completed the survey in Mandarin, Cantonese, or English. Household eligibility screening was performed for telephone and in-person interviews. The median response rate over 4 years of surveys was 21% based on Equation #3, the American Association for Public Opinion (AAPOR, 2011). A complete description of survey methods can be found elsewhere (Liao et al., 2011). A total of 3,001 respondents self-identified as Chinese American. Of those, 827 were aged 65 years. These analyses were limited to foreign-born respondents (n=805).

### Study Measures

Survey items were adapted from the Behavioral Risk Factor Surveillance System (BRFSS) interview, a previously validated survey (Nelson, Holtzman, Bolen, Stanwyck, & Mack, 2001).

## Dependent variables

**Health-related quality of life**—HRQOL is a health-focused quality of life measure of self-reported physical health, mental health, and social functioning (Centers for Disease Control, 2000) used to understand the health status of a population (Brown et al., 2003). *The Healthy People 2020* national health initiative includes health-related quality of life and well-being as one of its four overarching goals for all persons in the United States (U.S. Department of Health and Human Services, 2012). Four previously validated core HRQOL measures (Centers for Disease Control, 2000; Moriarty, Kobau, Zack, & Zahran, 2005; Moriarty, Zack, & Kobau, 2003) were used, including self-reported general health, days of poor physical health in the past month, days of poor mental health in the past month, and days of limited activities due to poor physical or mental health in the past month. General health was categorized as excellent/very good/good and fair/poor, while days of poor physical and mental health and limited activity days were categorized as  $\geq 14$  days and  $<14$  days in the past month.

## Behavioral Risk Factors

**Physical Activity**—PA levels were based on the *2008 Physical Activity Guidelines for Americans*, which recommends that all adults should perform  $\geq 150$  minutes of moderate intensity or  $\geq 75$  minutes of vigorous intensity PA weekly (U.S. Department of Health and Human Services, 2008). Respondents were classified as sufficiently active ( $\geq 150$  minutes/week of moderate PA,  $\geq 75$  minutes/week of vigorous PA, or an equivalent combination) or insufficiently active (less than sufficient or no PA).

**Cigarette Smoking**—Cigarette smoking status was determined using (1) “Have you smoked at least 100 cigarettes in your entire life?” and (2) “Do you now smoke cigarettes every day, some days, or not at all?” Those responding “yes” to the first question and “every day” or “some days” to the second question were current smokers. Those responding “no” to the first question were characterized as never smoked, and those responding “yes” to the first question and “not at all” to the second question were former smokers.

**Fruit and vegetable intake**—F&V intake was determined using a 6-item food frequency screener. Respondents reported consumption of 1) fruit juice; 2) fruit; 3) green salad; 4) potatoes (not counting fried potatoes or potato chips); 5) carrots; and 6) other vegetables. Healthy People 2010 objectives include consumption of  $\geq 3$  daily servings of vegetables and  $\geq 2$  daily servings of fruit (U.S. Department of Health and Human Services, 2000). For this analysis, frequencies were combined and adjusted to number of times per day and were dichotomized to  $\geq 5$  times per day versus  $<5$  times per day.

## Independent Variables

Socio-demographic variables included gender, age, education, and English language spoken at home. Body mass index (BMI) was calculated from self-reported height and weight. World Health Organization (WHO) recommended guidelines for Asians were adopted rather than the standard BMI definition, as a substantial proportion of Asian individuals with a high risk of type 2 diabetes have been found at lower BMI levels than the existing WHO

cut-off point of 25 for overweight (W.H.O. Expert Consultation, 2004). Asian BMI is defined as <23 (underweight/normal weight), 23-27.49 (overweight), and ≥27.5 (obese).

Additional variables included annual household income, health insurance, recent routine check-up, unable to see a doctor in the past year because of cost, diabetes diagnosis, high blood pressure diagnosis, leisure time PA, flu shot in the past year, and pneumonia vaccine in lifetime. English reading fluency was assessed in year 4.

## Statistical Analyses

Descriptive statistics were performed for all independent variables and stratified by gender. Percentages and standard errors (SEs) are presented. Chi-square tests were run for categorical variables and t-tests were run for continuous variables across gender.

Logistic regression models were run to identify the effects of gender, age, education, home language, and BMI on each outcome variable. Diagnoses of disease, variables with little variation such as health insurance, and variables found to be non-significant in bivariate analyses were not included in regression models. Income was excluded due to the large amount of missing responses. Adjusted odds ratios (ORs), 95% confidence intervals (CIs), and p-values were reported for all variables in each model. Two-sided p-values <0.05 were considered statistically significant. Hosmer and Lemeshow Goodness-of-Fit tests were conducted to test model fit.

SAS-callable SUDAAN 11.0 accounted for the complex sample design; each sample was weighted to reflect the probability of selection, the number of eligible family members, and the age-gender population size of the surveyed population.

## Results

Table 1 presents the socio-demographic and health characteristics of the sample, overall and by gender. Significant differences were found by gender. Women were more likely to have health insurance, have had a routine check-up in the past year, have been diagnosed with high blood pressure, and report more days of poor physical health, mental health, and limited activity. Men were more likely to have higher education, higher income, report better English reading fluency (year 4 only), smoke cigarettes, and perform recommended weekly PA.

Significant differences by gender were seen for all HRQOL measures; women self-reported worse health, in addition to a greater number of poor mental health, poor physical health, and limited activity days, when compared to men.

Overall, 66.8% of respondents reported insufficient levels of PA, 7.4% were current smokers, and 81.2% of respondents consumed F&V <5 times per day. Women were significantly more likely to receive insufficient PA and to be current non-smokers compared to men.

## HRQOL Measures

Results from a series of multivariable logistic regressions on HRQOL measures are presented in Table 2. For mental health, women were 2.07 times more likely than men to report 14 days of poor mental health in the past month. For physical health, women were 1.65 times more likely than men to report 14 days of poor physical health and older age was significantly associated with 14 days of poor physical health. For limited activity days, women were 2.06 times more likely than men to report 14 days of limited activities and older age was significantly associated with 14 days of limited activities. Compared to respondents who completed college, those with less than a high school education or some college education were significantly more likely to report fair or poor health (1.68 and 1.72 respectively). Finally, respondents with a BMI  $\geq 27.5$  were 1.80 times more likely than those with a BMI  $< 23$  to report fair or poor health.

## Behavioral Risk Factors

Results from a series of multivariable logistic regression on behavioral risk factors are presented in Table 3. Few demographic characteristics were significantly associated with the behavioral risk factors. Older age was significantly associated with insufficient PA. Due to the small smoking prevalence among women, adjusted analyses for smoking was only run for men. Men with less than a high school education were 6.13 times more likely than men with a college education to be current smokers and respondents with a BMI  $\geq 23$  were less likely than those with a BMI  $< 23$  to smoke.

## Discussion

To our knowledge, this is the first study to use a population-based dataset to oversample Asian Americans in NYC, providing important descriptive information on health behaviors and HRQOL among a large sample of older, foreign-born Chinese Americans. Men were significantly more likely than women to be current and former smokers. Only a third engaged in sufficient weekly PA, and men were significantly more likely than women to engage in sufficient PA. Among older Chinese respondents, HRQOL measures were generally low; the majority reported fair or poor health and days of poor health were high. Women were more likely to report poor mental health, poor physical health, and limited activity days than men.

Smoking prevalence (7.4%) was similar to overall smoking rates among older respondents in NYC (8.2%) (New York City Department of Health and Mental Hygiene, 2012); however, 14.8% of men and 1.3% of women were current smokers, and 33.8% of men and 3.8% of women were former smokers, similar to previous studies (Li, Kwon, Weerasinghe, Rey, & Trinh-Shevrin, 2013; Maxwell, Bernaards, & McCarthy, 2005; Yu, Chen, Kim, & Abdulrahim, 2002). As shown previously, lower educational attainment is associated with current smoking among men (Parikh et al., 2009). The finding that lower BMI is associated with cigarette smoking has been reported in a study among adults in China (Hou et al., 2008; Xu, Yin, & Wang, 2007), and studies have found that smokers often have lower body weight than nonsmokers (Chiolero, Faeh, Paccaud, & Cornuz, 2008; Flegal, Troiano, Pamuk, Kuczmarski, & Campbell, 1995). While this association is not completely understood, it



may be related to an increase in energy expenditure among smokers (Hofstetter, Schutz, Jequier, & Wahren, 1986).

Findings from national BRFSS data report that about half of adults aged 65 perform sufficient PA (Centers for Disease Control and Prevention, 2011). NYC 2012 data found that 38% of adults aged 65 performed sufficient PA and men are more likely than women to perform sufficient PA (New York City Department of Health and Mental Hygiene, 2012). A California study similarly reported that Asian Americans are less likely than non-Asians to meet recommended PA guidelines (August & Sorkin, 2011). As expected, older age was significantly associated with sedentary behaviors and receiving insufficient PA (DiPietro, 2001). Limitations to the PA measure do not allow us to determine reasons for lower PA rates in our sample. Potential areas for future research include the role of neighborhood factors such as perceptions of safety which have been found to influence PA engagement in older adults (Tucker-Seeley, Subramanian, Li, & Sorensen, 2009), as well as access to parks and walkability of neighborhoods (Wen, Kandula, & Lauderdale, 2007).

While recommended daily F&V intake was low, reported rates among NYC older adults and older Asian adults were lower (11.1% and 13.8%, respectively) (New York City Department of Health and Mental Hygiene, 2012). Research in California has shown low F&V intake among all adults; college-educated native- and foreign-born Asians had fewer servings of F&Vs than their less-educated counterparts (Bostean et al., 2013). Due to the limitations of the measures, reasons for low F&V intake among our sample are unclear.

In general, Asian Americans are less likely to report poorer HRQOL when compared to whites and other groups (Chowdhury, Balluz, & Strine, 2008; Zahran et al., 2005). Prior studies using 1993-2001 BRFSS data reported that about one-third of adults aged 65 self-reported fair or poor health (Zahran et al., 2005), whereas in our sample two-thirds of older adults self-reported fair or poor health. Women in our sample reported more days of poor physical and mental health than men, and similar findings have been shown among overall adults (McGuire et al., 2007; Zahran et al., 2005). Few studies have examined HRQOL specifically among older Asian populations; future studies should examine differences that may exist by ethnic subgroups.

These findings also underscore important study population characteristics. Approximately 62% reported having less than a high school education, similar to previous research on older NYC Chinese Americans (Parikh et al., 2009; Wong et al., 2013). In comparison, 26.9% of Chinese Americans aged 60 in California had less than a high school education (G. Kim, Chiriboga, et al., 2010). However, 89% of the older Chinese Americans from the California dataset were foreign-born and 56% had limited English proficiency (G. Kim, Chiriboga, et al., 2010). In contrast, our sample was all foreign-born, and in year 4, 89% of individuals reported reading English not well or not at all. Our findings are similar to NYC data, which reported that 88% of Chinese age 65 spoke English less than very well (Asian American Federation, 2013).

## Study Limitations

These findings have several limitations. The BRFSS F&V intake scale is less valid than other instruments; it may overestimate or underestimate daily intake (D. J. Kim & Holowaty, 2003). Second, key variables of interest, such as time in the U.S. and English reading fluency are missing or not included for all years. Research has found correlations between immigration age and health status (Lum & Vanderaa, 2010). Additionally, while health insurance was assessed, type of health insurance was not collected (e.g. private or public). Fourth, some predictor variables had high prevalence and could not be used in logistic regression, such as routine checkup and health insurance. Variables representing access to the healthcare system specific to older populations would be informative for these analyses.

## Implications for Practice

Findings suggest that this population differs from other groups of older adults, underscoring the need for further study. To address modifiable lifestyle behavioral risk factors and HRQOL among older Chinese Americans, strategies and policies should be culturally- and linguistically-tailored to ensure relevancy and successful uptake. Qualitative interviews among older Asians in NYC suggest the need to enhance access to social supports to improve the quality of life among older Asians coping with immigration-related stressors such as acculturation and language barriers and normal stresses of aging (Asian American Federation of New York, 2003). Yang & Kagawa-Singer propose that community-based, ethnic subgroup-specific healthcare organizations with bicultural and bilingual staff are needed to provide appropriate care to subgroups such as aging Chinese (Yang & Kagawa-Singer, 2007). Better understanding of cultural, generational, and linguistic challenges to mental health service utilization is critical in tailoring appropriate mental health and social service programs for this population (Asian American Federation of New York, 2003).

Health promotion programs should incorporate traditional cultures and health beliefs related to diet, as older adults may continue dietary patterns from their home country (Satia-Abouta, Patterson, Kristal, Teh, & Tu, 2002). Nutrition programs have focused on older adults with chronic diseases (Lyons, 2013); high rates of diabetes are shown among NYC Asian subgroups (Gupta, Wu, Young, & Perlman, 2011; Islam, Wyatt, et al., 2013), and several community-based, in-language, nutrition and chronic disease interventions have reportedly decreased chronic disease risk factors among targeted Chinese and other Asian Americans (Chesla et al., 2013; Islam, Zanolwiak, et al., 2013; M. T. Kim et al., 2009; Lv & Brown, 2011). Culturally relevant activities such as tai chi are also an effective way to promote PA and improve mental health (Adler & Roberts, 2006; Taylor-Piliae, Haskell, Stotts, & Froelicher, 2006; Wang, Collet, & Lau, 2004; Yeung et al., 2012). Community-based programs and system-level policies tailored to the social and cultural context of aging ethnic subpopulations are necessary to provide appropriate care and services to address chronic disease-related risk factors.

Important gender differences were noted and highlight the need for gender-specific, culturally-tailored interventions. While community-based and culturally-tailored interventions among Chinese American smokers in NYC and Pennsylvania have shown



positive results in improved self-efficacy and smoking cessation (Burton et al., 2010; Fang et al., 2006; Shelley et al., 2010; Wu et al., 2009), additional emphasis should be placed on older Chinese men. Furthermore, primary care-based interventions are needed to improve early detection of mental health concerns, especially among older Chinese women (Appel et al., 2011). Previous research found a significant association between income and poor mental health (Chung et al., 2003; Lubetkin, Jia, & Gold, 2003); the majority of our sample reported income earned less than \$25,000 annually, potentially placing them at greater risk. Studies have been proposed to address depression among Asian Americans (Trinh et al., 2011; Yeung, Hails, Chang, Trinh, & Fava, 2011), and a lifestyle intervention among older Chinese immigrants saw positive changes in mental health (Lu et al., 2014). Community-based, culturally-tailored interventions and policies that address factors associated with health-related behaviors and outcomes are key to providing services to the aging Chinese community as well as other aging racial and ethnic minority communities.

## Acknowledgments

### Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The NYU Center for the Study of Asian American Health is supported by the NIH/NIMHD cooperative agreement number 2P60MD000538; the NYU Health Promotion and Prevention Research Center is supported by the Centers for Disease Control and Prevention cooperative agreement number 1U48DP001904.

## References

- AAPOR. Standard Definitions: Final Dispositions of Case Codes and Outcome Rates for Surveys. 7th edition. 2011. Retrieved from <http://aapor.org/Content/NavigationMenu/AboutAAPOR/StandardsandEthics/StandardDefinitions/StandardDefinitions2011.pdf>
- Adler PA, Roberts BL. The use of Tai Chi to improve health in older adults. *Orthopaedic Nursing*. 2006; 25(2):122–126. [PubMed: 16572030]
- Alegria M, Chatterji P, Wells K, Cao Z, Chen CN, Takeuchi D, Meng XL. Disparity in depression treatment among racial and ethnic minority populations in the United States. *Psychiatric Services*. 2008; 59(11):1264–1272. [PubMed: 18971402]
- Appel HB, Huang B, Ai AL, Lin CJ. Physical, behavioral, and mental health issues in Asian American women: results from the National Latino Asian American Study. *Journal of Women's Health*. 2011; 20(11):1703–1711.
- Asian American Federation. Profile of New York City's Chinese Americans. 2013. p. 2013 Retrieved from <http://www.aafny.org/cic/briefs/chinese2013.pdf>
- Asian American Federation of New York. Asian American elders in New York City: A study of health, social needs, quality of life and quality of care. 2003. Retrieved from <http://www.aafny.org/doc/AsianAmericanEldersInNYC.pdf>
- August KJ, Sorkin DH. Racial/ethnic disparities in exercise and dietary behaviors of middle-aged and older adults. *Journal of General Internal Medicine*. 2011; 26(3):245–250. [PubMed: 20865342]
- Bostean G, Roberts CK, Crespi CM, Prelip M, Peters A, Belin TR, McCarthy WJ. Cardiovascular health: associations with race-ethnicity, nativity, and education in a diverse, population-based sample of Californians. *Annals of epidemiology*. 2013; 23(7):388–394. [PubMed: 23726820]
- Brown DW, Balluz LS, Heath GW, Moriarty DG, Ford ES, Giles WH, Mokdad AH. Associations between recommended levels of physical activity and health-related quality of life. Findings from the 2001 Behavioral Risk Factor Surveillance System (BRFSS) survey. *Preventive Medicine*. 2003; 37(5):520–528. [PubMed: 14572437]

- Burton D, Zeng XX, Chiu CH, Sun J, Sze NL, Chen Y, Chin MS. A phone-counseling smoking-cessation intervention for male Chinese restaurant workers. *Journal of Community Health*. 2010; 35(6):579–585. [PubMed: 20186474]
- Buttar HS, Li T, Ravi N. Prevention of cardiovascular diseases: Role of exercise, dietary interventions, obesity and smoking cessation. *Experimental & Clinical Cardiology*. 2005; 10(4):229–249. [PubMed: 19641674]
- Carlson SA, Fulton JE, Schoenborn CA, Loustalot F. Trend and prevalence estimates based on the 2008 Physical Activity Guidelines for Americans. *American Journal of Preventive Medicine*. 2010; 39(4):305–313. [PubMed: 20837280]
- Centers for Disease Control. Measuring healthy days: Population Assessment of Health-Related Quality of Life. 2000. Retrieved from <http://www.cdc.gov/hrqol/pdfs/mhd.pdf>
- Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System: Prevalence and Trends Data. 2011. Retrieved from <http://apps.nccd.cdc.gov/brfss/>
- Chesla CA, Chun KM, Kwan CM, Mullan JT, Kwong Y, Hsu L, Waters CM. Testing the efficacy of culturally adapted coping skills training for Chinese American immigrants with type 2 diabetes using community-based participatory research. *Research in Nursing & Health*. 2013; 36(4):359–372. [PubMed: 23606271]
- Chiolero A, Faeh D, Paccaud F, Cornuz J. Consequences of smoking for body weight, body fat distribution, and insulin resistance. *The American journal of clinical nutrition*. 2008; 87(4):801–809. [PubMed: 18400700]
- Chowdhury PP, Balluz L, Strine TW. Health-related quality of life among minority populations in the United States, BRFSS 2001–2002. *Ethnicity & Disease*. 2008; 18(4):483–487. [PubMed: 19157254]
- Chung H, Teresi J, Guarnaccia P, Meyers BS, Holmes D, Bobrowitz T, Ferran E Jr. Depressive symptoms and psychiatric distress in low income Asian and Latino primary care patients: prevalence and recognition. *Community Mental Health Journal*. 2003; 39(1):33–46. [PubMed: 12650554]
- Danaei G, Ding EL, Mozaffarian D, Taylor B, Rehm J, Murray CJ, Ezzati M. The preventable causes of death in the United States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors. *PLoS medicine*. 2009; 6(4):e1000058. [PubMed: 19399161]
- DiPietro L. Physical activity in aging: changes in patterns and their relationship to health and function. *The Journals of Gerontology, Series A: Biological Sciences and Medical Sciences*. 2001; 56:13–22. Spec No 2.
- Drewnowski A, Evans WJ. Nutrition, physical activity, and quality of life in older adults: summary. *The Journals of Gerontology, Series A: Biological Sciences and Medical Sciences*. 2001; 56:89–94. Spec No 2.
- Fang CY, Ma GX, Miller SM, Tan Y, Su X, Shive S. A brief smoking cessation intervention for Chinese and Korean American smokers. *Preventive Medicine*. 2006; 43(4):321–324. [PubMed: 16860858]
- Flegal KM, Troiano RP, Pamuk ER, Kuczmarski RJ, Campbell SM. The influence of smoking cessation on the prevalence of overweight in the United States. *The New England journal of medicine*. 1995; 333(18):1165–1170. [PubMed: 7565970]
- Gregg EW, Pereira MA, Caspersen CJ. Physical activity, falls, and fractures among older adults: a review of the epidemiologic evidence. *Journal of the American Geriatrics Society*. 2000; 48(8):883–893. [PubMed: 10968291]
- Gupta LS, Wu CC, Young S, Perlman SE. Prevalence of diabetes in New York City, 2002–2008: comparing foreign-born South Asians and other Asians with U.S.-born whites, blacks, and Hispanics. *Diabetes Care*. 2011; 34(8):1791–1793. [PubMed: 21715521]
- Haveman-Nies A, de Groot LC, van Staveren WA. Dietary quality, lifestyle factors and healthy ageing in Europe: the SENECA study. *Age and ageing*. 2003; 32(4):427–434. [PubMed: 12851188]
- He, W.; Census Bureau, US. Current Population Reports, Series P23-211, The Older Foreign-Born Population of the United States. U.S. Government Printing Office; Washington, D.C.: 2002. p. 2000

- Hofstetter A, Schutz Y, Jequier E, Wahren J. Increased 24-hour energy expenditure in cigarette smokers. *The New England journal of medicine*. 1986; 314(2):79–82. [PubMed: 3941694]
- Hou X, Jia W, Bao Y, Lu H, Jiang S, Zuo Y, Xiang K. Risk factors for overweight and obesity, and changes in body mass index of Chinese adults in Shanghai. *BMC Public Health*. 2008; 8:389. [PubMed: 19025585]
- Hwang WC, Chun CA, Takeuchi DT, Myers HF, Siddarth P. Age of first onset major depression in Chinese Americans. *Cultural Diversity & Ethnic Minority Psychology*. 2005; 11(1):16–27. [PubMed: 15727492]
- Islam NS, Khan S, Kwon S, Jang D, Ro M, Trinh-Shevrin C. Methodological issues in the collection, analysis, and reporting of granular data in Asian American populations: historical challenges and potential solutions. *Journal of Health Care for the Poor and Underserved*. 2010; 21(4):1354–1381. [PubMed: 21099084]
- Islam NS, Wyatt LC, Kapadia SB, Rey MJ, Trinh-Shevrin C, Kwon SC. Diabetes and associated risk factors among Asian American subgroups in New York City. *Diabetes Care*. 2013; 36(1):e5. [PubMed: 23264307]
- Islam NS, Zanolwaki JM, Wyatt LC, Chun K, Lee L, Kwon SC, Trinh-Shevrin C. A Randomized-Controlled, Pilot Intervention on Diabetes Prevention and Healthy Lifestyles in the New York City Korean Community. *Journal of Community Health*. 2013; 38(6):1030–1041. [PubMed: 23813322]
- Kim DJ, Holowaty EJ. Brief, validated survey instruments for the measurement of fruit and vegetable intakes in adults: a review. *Preventive Medicine*. 2003; 36(4):440–447. [PubMed: 12649052]
- Kim G, Chiriboga DA, Jang Y, Lee S, Huang CH, Parmelee P. Health status of older Asian Americans in California. *Journal of the American Geriatrics Society*. 2010; 58(10):2003–2008. [PubMed: 20929469]
- Kim G, Jang Y, Chiriboga DA, Ma GX, Schonfeld L. Factors associated with mental health service use in Latino and Asian immigrant elders. *Aging & Mental Health*. 2010; 14(5):535–542. [PubMed: 20496182]
- Kim MT, Han HR, Song HJ, Lee JE, Kim J, Ryu JP, Kim KB. A community-based, culturally tailored behavioral intervention for Korean Americans with type 2 diabetes. *The Diabetes Educator*. 2009; 35(6):986–994. [PubMed: 19934458]
- Lau DS, Lee G, Wong CC, Fung GL, Cooper BA, Mason DT. Characterization of systemic hypertension in the San Francisco Chinese community. *American Journal of Cardiology*. 2005; 96(4):570–573. [PubMed: 16098313]
- Li S, Kwon SC, Weerasinghe I, Rey MJ, Trinh-Shevrin C. Smoking among Asian Americans: acculturation and gender in the context of tobacco control policies in New York City. *Health Promotion Practice*. 2013; 14(5 Suppl):18S–28S. [PubMed: 23667057]
- Liao Y, Bang D, Cosgrove S, Dulin R, Harris Z, Taylor A, Centers for Disease Control and Prevention. Surveillance of health status in minority communities - Racial and Ethnic Approaches to Community Health Across the U.S. (REACH U.S.) Risk Factor Survey, United States, 2009. *Morbidity and Mortality Weekly Report: Surveillance Summaries*. 2011; 60(6):1–44. [PubMed: 21597458]
- Lloyd-Jones DM, Leip EP, Larson MG, D'Agostino RB, Beiser A, Wilson PW, Levy D. Prediction of lifetime risk for cardiovascular disease by risk factor burden at 50 years of age. *Circulation*. 2006; 113(6):791–798. [PubMed: 16461820]
- Lu Y, Dipierro M, Chen L, Chin R, Fava M, Yeung A. The evaluation of a culturally appropriate, community-based lifestyle intervention program for elderly Chinese immigrants with chronic diseases: a pilot study. *Journal of Public Health*. 2014; 36(1):149–155. [PubMed: 23596196]
- Lubetkin EI, Jia H, Gold MR. Depression, anxiety, and associated health status in low-income Chinese patients. *American Journal of Preventive Medicine*. 2003; 24(4):354–360. [PubMed: 12726874]
- Lum TY, Vanderka JP. Health disparities among immigrant and non-immigrant elders: the association of acculturation and education. *Journal of Immigrant and Minority Health*. 2010; 12(5):743–753. [PubMed: 19184599]
- Lv N, Brown JL. Impact of a nutrition education program to increase intake of calcium-rich foods by Chinese-American women. *Journal of the American Dietetic Association*. 2011; 111(1):143–149. [PubMed: 21185977]

- Lyons BP. Nutrition Education Intervention with Community-Dwelling Older Adults: Research Challenges and Opportunities. *Journal of Community Health*(Advance online publication. 2013 doi:10.1007/s10900-013-9810-x).
- Maxwell AE, Bernaards CA, McCarthy WJ. Smoking prevalence and correlates among Chinese- and Filipino-American adults: findings from the 2001 California Health Interview Survey. *Preventive Medicine*. 2005; 41(2):693–699. [PubMed: 15917070]
- McGuire LC, Strine TW, Okoro CA, Ahluwalia IB, Ford ES. Modifiable characteristics of a healthy lifestyle in U.S. older adults with or without frequent mental distress: 2003 Behavioral Risk Factor Surveillance System. *The American Journal of Geriatric Psychiatry*. 2007; 15(9):754–761. [PubMed: 17804829]
- Moriarty DG, Kobau R, Zack MM, Zahran HS. Tracking Healthy Days -- a window on the health of older adults. *Preventing Chronic Disease*. 2005; 2(3):A16. [PubMed: 15963318]
- Moriarty DG, Zack MM, Kobau R. The Centers for Disease Control and Prevention's Healthy Days Measures - population tracking of perceived physical and mental health over time. *Health and Quality of Life Outcomes*. 2003; 1:37. [PubMed: 14498988]
- Mozaffarian D, Afshin A, Benowitz NL, Bittner V, Daniels SR, Franch HA, Metabolism C. o. C. C. C. o. C. D. i. t. Y. C. o. t. K. i. C. Population approaches to improve diet, physical activity, and smoking habits: a scientific statement from the American Heart Association. *Circulation*. 2012; 126(12):1514–1563. [PubMed: 22907934]
- Nelson DE, Holtzman D, Bolen J, Stanwyck CA, Mack KA. Reliability and validity of measures from the Behavioral Risk Factor Surveillance System (BRFSS). *Social and Preventive Medicine*. 2001; 46(Suppl 1):S3–42. [PubMed: 11851091]
- New York City Department of Health and Mental Hygiene. Epiquery: NYC Interactive Health Data System - Community Health Survey. 2012. Retrieved from <http://nyc.gov/health/epiquery>
- Paganini-Hill A. Lifestyle practices and cardiovascular disease mortality in the elderly: the leisure world cohort study. *Cardiology Research and Practice*. 2011; 2011 Article ID 983764, 7 pages, doi:10.4061/2011/983764.
- Parikh NS, Fahs MC, Shelley D, Yerneni R. Health behaviors of older Chinese adults living in New York City. *Journal of Community Health*. 2009; 34(1):6–15. [PubMed: 18830807]
- Physical Activity Guidelines Advisory Committee. Physical Activity Guidelines Advisory Committee Report. 2008. Retrieved from <http://www.health.gov/paguidelines/committeereport.aspx>
- Pollack H, Wang S, Wyatt L, Peng CH, Wan K, Trinh-Shevrin C, Kwon S. A comprehensive screening and treatment model for reducing disparities in hepatitis B. *Health Affairs*. 2011; 30(10):1974–1983. [PubMed: 21976342]
- Rajpathak SN, Wylie-Rosett J. High prevalence of diabetes and impaired fasting glucose among Chinese immigrants in New York City. *Journal of Immigrant and Minority Health*. 2011; 13(1):181–183. [PubMed: 20533090]
- Satia-Abouta J, Patterson RE, Kristal AR, Teh C, Tu SP. Psychosocial predictors of diet and acculturation in Chinese American and Chinese Canadian women. *Ethnicity & health*. 2002; 7(1):21–39. [PubMed: 12119064]
- Shah AM, Guo L, Magee M, Cheung W, Simon M, LaBrecche A, Liu H. Comparing selected measures of health outcomes and health-seeking behaviors in Chinese, Cambodian, and Vietnamese communities of Chicago: results from local health surveys. *Journal of Urban Health*. 2010; 87(5):813–826. [PubMed: 20625846]
- Shelley D, Nguyen N, Peng CH, Chin M, Chang MD, Fahs M. Increasing access to evidence-based smoking cessation treatment: effectiveness of a free nicotine patch program among Chinese immigrants. *Journal of Immigrant and Minority Health*. 2010; 12(2):198–205. [PubMed: 18825498]
- Strawbridge WJ, Deleger S, Roberts RE, Kaplan GA. Physical activity reduces the risk of subsequent depression for older adults. *American Journal of Epidemiology*. 2002; 156(4):328–334. [PubMed: 12181102]

- Takeuchi DT, Zane N, Hong S, Chae DH, Gong F, Gee GC, Alegria M. Immigration-related factors and mental disorders among Asian Americans. *American Journal of Public Health*. 2007; 97(1): 84–90. [PubMed: 17138908]
- Taylor-Piliae RE, Haskell WL, Stotts NA, Froelicher ES. Improvement in balance, strength, and flexibility after 12 weeks of Tai chi exercise in ethnic Chinese adults with cardiovascular disease risk factors. *Alternative Therapies in Health and Medicine*. 2006; 12(2):50–58. [PubMed: 16541997]
- Trinh-Shevrin, C.; Islam, NS.; Rey, MJ. *Asian American Communities and Health*. Jossey-Bass; San Francisco, CA: 2009.
- Trinh-Shevrin C, Pollack HJ, Tsang T, Park J, Ramos MR, Islam N, Kwon SC. The Asian American hepatitis B program: building a coalition to address hepatitis B health disparities. *Progress in Community Health Partnerships*. 2011; 5(3):261–271. [PubMed: 22080774]
- Trinh NH, Bedoya CA, Chang TE, Flaherty K, Fava M, Yeung A. A study of a culturally focused psychiatric consultation service for Asian American and Latino American primary care patients with depression. *BMC Psychiatry*. 2011; 11:166. [PubMed: 21995514]
- Tucker-Seeley RD, Subramanian SV, Li Y, Sorensen G. Neighborhood safety, socioeconomic status, and physical activity in older adults. *American Journal of Preventive Medicine*. 2009; 37(3):207–213. [PubMed: 19595554]
- U.S. Census Bureau. Profile of General Demographic Characteristics: 2000. 2000. Retrieved from <http://factfinder2.census.gov/>
- U.S. Census Bureau. Profile of the General Population and Housing Characteristics: 2010. 2010. Retrieved from <http://factfinder2.census.gov/>
- U.S. Department of Health and Human Services. Healthy People 2010. 2000. Retrieved from <http://www.healthypeople.gov/>
- U.S. Department of Health and Human Services. 2008 Physical Activity Guidelines for Americans. 2008. Retrieved from <http://www.health.gov/paguidelines/pdf/paguide.pdf>
- U.S. Department of Health and Human Services. The Vision, Mission, and Goals of Healthy People 2020. 2012. Retrieved from <http://healthypeople.gov/2020/Consortium/HP2020Framework.pdf>
- W.H.O. Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*. 2004; 363(9403):157–163. [PubMed: 14726171]
- Wang C, Collet JP, Lau J. The effect of Tai Chi on health outcomes in patients with chronic conditions: a systematic review. *Archives of Internal Medicine*. 2004; 164(5):493–501. [PubMed: 15006825]
- Wen M, Kandula NR, Lauderdale DS. Walking for transportation or leisure: what difference does the neighborhood make? *Journal of General Internal Medicine*. 2007; 22(12):1674–1680. [PubMed: 17932724]
- Wong SS, Dixon LB, Gilbride JA, Kwan TW, Stein RA. Measures of acculturation are associated with cardiovascular disease risk factors, dietary intakes, and physical activity in older Chinese Americans in New York City. *Journal of Immigrant and Minority Health*. 2013; 15(3):560–568. [PubMed: 22752686]
- Wu D, Ma GX, Zhou K, Zhou D, Liu A, Poon AN. The effect of a culturally tailored smoking cessation for Chinese American smokers. *Nicotine & Tobacco Research*. 2009; 11(12):1448–1457. [PubMed: 19915080]
- Xu F, Yin XM, Wang Y. The association between amount of cigarettes smoked and overweight, central obesity among Chinese adults in Nanjing, China. *Asia Pacific Journal of Clinical Nutrition*. 2007; 16(2):240–247. [PubMed: 17468078]
- Yang JS, Kagawa-Singer M. Increasing access to care for cultural and linguistic minorities: ethnicity-specific health care organizations and infrastructure. *Journal of Health Care for the Poor and Underserved*. 2007; 18(3):532–549. [PubMed: 17675712]
- Yeung A, Chan R, Mischoulon D, Sonawalla S, Wong E, Nierenberg AA, Fava M. Prevalence of major depressive disorder among Chinese-Americans in primary care. *General Hospital Psychiatry*. 2004; 26(1):24–30. [PubMed: 14757299]

- Yeung A, Hails K, Chang T, Trinh NH, Fava M. A study of the effectiveness of telepsychiatry-based culturally sensitive collaborative treatment of depressed Chinese Americans. *BMC Psychiatry*. 2011; 11:154. [PubMed: 21943315]
- Yeung A, Lepoutre V, Wayne P, Yeh G, Slipp LE, Fava M, Fricchione GL. Tai chi treatment for depression in Chinese Americans: a pilot study. *American Journal of Physical Medicine & Rehabilitation*. 2012; 91(10):863–870. [PubMed: 22790795]
- Yeung A, Yu SC, Fung F, Vorono S, Fava M. Recognizing and engaging depressed Chinese Americans in treatment in a primary care setting. *International Journal of Geriatric Psychiatry*. 2006; 21(9):819–823. [PubMed: 16955440]
- Yu ES, Chen EH, Kim KK, Abdulrahim S. Smoking among Chinese Americans: behavior, knowledge, and beliefs. *American Journal of Public Health*. 2002; 92(6):1007–1012. [PubMed: 12036797]
- Zahrn HS, Kobau R, Moriarty DG, Zack MM, Holt J, Donehoo R, Centers for Disease Control and Prevention. Health-related quality of life surveillance--United States, 1993-2002. *Morbidity and Mortality Weekly Report: Surveillance Summaries*. 2005; 54(4):1–35. [PubMed: 16251867]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Table 1**

Sociodemographics and Health-Related Characteristics of Foreign-Born Chinese Individuals Aged 65 in New York City by Gender and Percentage (SE).

Characteristics	Total (n=805)	Men (n=392)	Women (n=413)	p value
<i>Sociodemographic characteristics</i>				
Age (years)				0.001
65 to 69	26.0 (1.7)	25.9 (2.4)	26.2 (2.3)	
70 to 74	26.7 (1.7)	24.5 (2.3)	28.6 (2.3)	
75 to 79	23.1 (1.6)	28.9 (2.3)	18.0 (2.0)	
80 and over	24.2 (1.6)	20.7 (2.0)	27.2 (2.2)	
Education				0.007
Less than high school	62.4 (1.9)	57.4 (2.6)	66.8 (2.5)	
High school graduate	23.6 (1.6)	25.2 (2.3)	22.2 (2.1)	
College graduate	14.0 (1.3)	17.4 (1.9)	11.0 (1.6)	
Income				0.004
<\$25,000	73.5 (1.8)	72.6 (2.4)	74.2 (2.3)	
\$25,000	16.2 (1.5)	19.9 (2.1)	13.0 (1.7)	
Missing/Don't know	10.3 (1.2)	7.5 (1.4)	12.8 (1.8)	
English spoken at home	18.6 (1.6)	18.8 (2.1)	18.5 (2.0)	0.895
How well do you read English (2012 only, n=187)				0.030
Very well	1.9 (0.9)	1.0 (1.0)	3.2 (1.8)	
Well	9.3 (2.3)	12.8 (3.3)	4.2 (2.1)	
Not well	40.0 (3.7)	43.4 (5.1)	35.3 (5.1)	
Not at all	48.8 (3.9)	42.9 (5.2)	57.4 (5.2)	
<i>Health care usage and diagnoses</i>				
Health Care Coverage	97.1 (0.6)	95.4 (1.1)	98.6 (0.6)	0.009
Check-up in the last year	91.7 (1.0)	89.4 (1.6)	93.7 (1.3)	0.037
Too costly to see a doctor in the past year	7.0 (1.0)	8.1 (1.5)	6.0 (1.2)	0.251
BMI, mean (SE)	24.0 (0.2)	24.1 (0.2)	23.9 (0.2)	0.426
BMI (Asian definition)				0.080
Underweight/Normal weight (<23 kg/m <sup>2</sup> )	41.4 (1.9)	37.7 (2.6)	44.7 (2.6)	
Overweight (23-27.49 kg/m <sup>2</sup> )	44.5 (1.9)	49.0 (2.7)	40.7 (2.6)	
Obese (≥ 27.50 kg/m <sup>2</sup> )	14.1 (1.3)	13.4 (1.8)	14.6 (1.8)	
Diabetes diagnosis	27.5 (1.7)	26.0 (2.3)	28.7 (2.3)	0.405
High blood pressure diagnosis	58.5 (1.8)	52.9 (2.7)	63.2 (2.5)	0.005
Flu shot in past 12 months	85.5 (1.4)	82.9 (2.0)	87.7 (1.7)	0.058
Pneumonia vaccination in lifetime	46.8 (2.0)	46.0 (2.6)	47.4 (2.6)	0.277
Health behaviors				
Cigarette Smoking				<0.001
Current smoker	7.4 (1.0)	14.8 (2.0)	1.3 (0.6)	
Former smoker	17.4 (1.4)	33.8 (2.5)	3.8 (0.9)	

Characteristics	Total (n=805)	Men (n=392)	Women (n=413)	p value
Non-smoker	75.2 (1.6)	51.4 (2.7)	95.0 (1.1)	
Any PA in past month	70.6 (1.7)	72.3 (2.4)	69.3 (2.4)	0.367
Recommended PA				0.020
None/Insufficient	66.8 (1.8)	62.5 (2.5)	70.4 (2.4)	
At least sufficient	33.2 (1.8)	37.5 (2.5)	29.6 (2.4)	
Daily F&V Intake, mean (SE)	3.6 (0.1)	3.6 (0.1)	3.6 (0.1)	0.643
<3 times	35.5 (1.9)	36.7 (2.5)	34.6 (2.5)	0.778
3 to <5 times	45.6 (1.9)	45.3 (2.7)	45.9 (2.6)	
5 times	18.8 (1.5)	18.0 (2.0)	19.5 (2.1)	
Health-Related Quality of Life				
Self-rated health status				0.106
Excellent/very good/good	38.3 (1.9)	41.3 (2.6)	35.8 (2.5)	
Fair/poor	61.7 (1.9)	58.7 (2.6)	64.2 (2.5)	
Poor physical health days in past month, mean (SE)	5.3 (0.4)	4.4 (0.5)	6.1 (0.5)	0.016
14 days	16.0 (1.4)	13.1 (1.8)	18.6 (2.1)	0.009
1 to 13 days	24.9 (1.7)	21.9 (2.2)	27.5 (2.4)	
No days	59.1 (2.0)	65.0 (2.6)	54.0 (2.7)	
Poor mental health days in past month, mean (SE)	2.6 (0.3)	1.9 (0.3)	3.3 (0.4)	0.004
14 days	8.0 (1.0)	5.3 (1.2)	10.4 (1.7)	0.013
1 to 13 days	16.3 (1.5)	14.5 (1.8)	17.9 (2.1)	
No days	75.7 (1.7)	80.2 (2.1)	71.7 (2.4)	
Limited activity days in past month, mean (SE)	2.3 (0.3)	1.6 (0.3)	2.8 (0.4)	0.010
14 days	6.1 (0.9)	4.1 (1.0)	7.9 (1.4)	0.029
1 to 13 days	12.3 (1.3)	10.7 (1.6)	13.7 (1.8)	
No days	81.6 (1.5)	85.3 (1.8)	78.4 (2.2)	

Note. BMI = body mass index; PA = physical activity; F&V = fruit and vegetable

Table 2

Predictors of Health-Related Quality of Life Measures Among Foreign-Born Chinese Individuals Aged 65, Adjusted Logistic Regression Models

Predictor	14 days of poor mental health (Ref=<14 days)	p value	14 days of poor physical health (Ref=<14days)	p value	14 days of limited activities (Ref=<14 days)	p value	Fair/Poor SRHS (Ref = good or higher)	p value
Gender (ref=male)								
Female	2.07 (1.12, 3.84)	0.021	1.65 (1.07, 2.53)	0.022	2.06 (1.03, 4.11)	0.041	1.23 (0.91, 1.66)	0.185
Age	1.04 (0.99, 1.08)	0.122	1.05 (1.02, 1.09)	<0.001	1.10 (1.04, 1.16)	0.001	1.01 (0.98, 1.03)	0.513
Education (ref=college graduate)								
Less than high school	0.62 (0.26, 1.51)	0.292	0.73 (0.40, 1.35)	0.319	1.22 (0.44, 3.36)	0.703	1.68 (1.05, 2.68)	0.031
High school/some college	1.10 (0.40, 3.00)	0.854	0.71 (0.35, 1.45)	0.344	0.59 (0.17, 2.08)	0.413	1.72 (1.03, 2.87)	0.037
Home language (ref=English not spoken)								
English spoken	1.06 (0.48, 2.33)	0.893	1.47 (0.86, 2.50)	0.159	0.69 (0.27, 1.79)	0.446	1.13 (0.73, 1.75)	0.585
BMI (ref = Underweight/Normal <23)								
Overweight (23-27.49)	1.14 (0.59, 2.20)	0.686	0.97 (0.61, 1.54)	0.888	1.41 (0.69, 2.88)	0.345	1.06 (0.75, 1.49)	0.750
Obese ( ≥ 27.5)	1.86 (0.84, 4.11)	0.126	1.12 (0.60, 2.10)	0.712	1.13 (0.38, 3.36)	0.826	1.80 (1.09, 2.96)	0.021

Note. SRHS = self-rated health status; BMI = body mass index

**Table 3**  
Predictors of Behavioral Risk Factors among Foreign-Born Chinese Individuals Aged 65, adjusted logistic regression models

Predictor	Less Than Sufficient Exercise (Ref= Sufficient Exercise)	p value	Current Smoker, Males Only (Ref=Nonsmoker)	p value	Fruits & Vegetables <5 Times/Day (Ref= 5 Times/Day)	p value
Gender (ref=male)						
Female	1.34 (0.97, 1.85)	0.073	N/A		0.80 (0.55, 1.17)	0.251
Age	1.05 (1.02, 1.85)	<0.001	0.95 (0.90, 1.00)	0.064	1.00 (0.97, 1.03)	0.829
Education (ref=College graduate)						
Less than high school	1.46 (0.92, 2.30)	0.104	6.13 (1.82, 20.63)	0.003	1.60 (0.94, 2.74)	0.086
High school/some college	1.05 (0.64, 1.73)	0.844	2.86 (0.74, 11.08)	0.128	1.46 (0.80, 2.68)	0.222
Home language (ref=English not spoken)						
English spoken	1.09 (0.71, 1.67)	0.688	1.14 (0.49, 2.65)	0.761	0.95 (0.56, 1.59)	0.842
BMI (ref = Underweight/Normal <23)						
Overweight (23-27.49)	0.88 (0.61, 1.27)	0.502	0.36 (0.17, 0.73)	0.005	0.71 (0.47, 1.08)	0.107
Obese ( ≥ 27.5)	1.14 (0.68, 1.91)	0.609	0.26 (0.09, 0.76)	0.014	0.90 (0.49, 1.65)	0.740

Note. BMI = body mass index